# Stream Insects and Crustaceans ID Card

Lines under picture indicate the relative size of organisms



# Aquatic Worm: Class Oligocheata

½" - 2", can be very tiny; thin, wormlike body, tolerant of impairment



#### Flat Worm:

#### Family Planaridae


Up to  $\frac{1}{4}$ ", soft body, may have distinct head with eyespots, tolerant of impairment



#### Leech:

#### Order Hirudinea

 $\frac{1}{4}$ " - 2", segmented body, suction cups on both ends, tolerant of impairment



#### Crayfish: Order Decapoda

Up to 6", 2 large claws, 8 legs, resembles a small lobster, somewhat tolerant of impairment



## Sowbug: Order Isopoda

 $\frac{1}{4}$ " -  $\frac{3}{4}$ ", gray oblong body wider than it is high, more than 6 legs, long antennae, somewhat tolerant of impairment



#### Scud: Order Amphipoda

½", white to gray, body higher than it is wide, swims sideways, more than 6 legs, resembles small shrimp, somewhat tolerant of impairment



#### Stonefly: Order Plecoptera

 $\frac{1}{2}$ " - 1  $\frac{1}{2}$ ", 6 legs with hooked tips, antennae, 2 hair-like tails, no gills on abdomen, very intolerant of impairment



### Mayfly:

### Order Emphemeroptera

 $\frac{1}{4}$ " - 1", plate-like or feathery gills on abdomen, 6 hooked legs, 2 or 3 long hair-like tails, tails may be webbed together, very intolerant of impairment



# Dragonfly and Damselfly: Order Odonata

 $\frac{1}{2}$  " - 2", large eyes, 6 hooked legs, large protracting lower jaw, 3 broad oar-shaped tails OR wide oval to round abdomen, somewhat tolerant of impairment



#### Hellgrammite, Fishfly, and Alderfly: Order Megaloptera

 $\frac{3}{4}$ " - 4", 6 legs, large pinching jaws, 8 pairs of feelers along abdomen, 2 hooks on tail end OR 1 single spiky tail, somewhat tolerant of impairment



#### Common Netspinners: Family Hydropsychidae

Up to  $\frac{3}{4}$ ", 6 hooked legs on upper 1/3 of body, 2 hooks at back end, underside of abdomen with white tufts of gills, somewhat tolerant of impairment



#### Most Caddisfly: Order Trichoptera

Up to 1", 6 hooked legs on upper 1/3 of body, may be in stick, rock or leaf case, no gill tufts on abdomen, intolerant of impairment

Illustrations from: Voshell, J. R., Jr. 2001. Guide to the Common Freshwater Invertebrates of North America. MacDonald and Woodward Publishing Co. With permission of the author.

# Stream Insects and Crustaceans ID Card

Lines under picture indicate the relative size of organisms



#### Beetles: Order Coleoptera

 $\frac{1}{4}$ " - 1", disk-like oval body with 6 small legs and gill tufts on underside OR small black beetle crawling on streambed OR commalike brown "crunchy" body with 6 legs on upper 1/3 and possibly gill tuft on back end, OR (miscellaneous body form - rare), somewhat tolerant of impairment



#### Midges:


#### Family Chironomidae

Up to  $\frac{1}{4}$ ", distinct head, worm-like segmented body, 2 leg-like projections on each side, often whitish to clear, occasionally bright red, tolerant of impairment



#### Black Fly: Family Simuliidae

Up to  $\frac{1}{4}$ ", end of body wider (like bowling pin), distinctive head, sucker on end, tolerant of impairment



#### Most True Flies: Order Diptera

 $\frac{1}{4}$ " - 2", bodies plump and maggotlike, may have caterpillar like "legs" along body, may have lobes or conical tails on end, tolerant of impairment



#### Gilled Snails: Class Gastropoda

Up to  $\frac{3}{4}$ ", shell opening covered by a thin plate called an operculum, with helix pointed up shell opens to the right, intolerant of impairment



# Lunged Snails: Class Gastropoda

Up to  $\frac{3}{4}$ ", no operculum, with helix pointed up shell opens to the left, tolerant of impairment




#### Clams: Class Bivalvia

Up to  $\frac{3}{4}$ ", fleshy body enclosed between two clamped together shells (if clam is alive, shells cannot be pried apart without harming clam), somewhat tolerant of impairment



#### Head



Tails: There are many different kinds of macroinvertebrate tails. The thin thread-like tails found on stoneflies and mayflies are called cerci. The oar-shaped tails found on a damselfly are not really tails they are actually gills called caudal lamellae!



#### VA Save Our Streams Program

VA Division of the Izaak Walton League of America P.O. Box 8297 Richmond, VA 23226 (804) 615-5036 www.vasos.org

These sheets are modified from the National Izaak Walton League of America SOS Program Stream Insects & Crustaceans ID Card.

http://www.iwla.org/SOS/index.html

Illustrations from: Voshell, J. R., Jr. 2001. Guide to the Common Freshwater Invertebrates of North America. MacDonald and Woodward Publishing Co. With permission of the author.